COMPARISONS OF PREOPERATIVE BLOOD PRESSURES IN SURGICAL PATIENTS

Brian Stacy BSN, RN; Kenneth Daratha PhD Gonzaga University and Providence Sacred Heart Medical Center

Background

Surgical patients expect anesthesia providers to provide high quality care. Blood pressure variability must be mitigated by establishing the patient's blood pressure baseline and sustaining it through surgery. A consensus has not been established on which blood pressure measurement should be used as the patient's baseline.

Blood pressures before surgery in multiple care settings have been shown to vary. This may result in differences in drug and fluid administration, depending on the blood pressure considered baseline by the anesthesia provider.

The purpose of this observational evidence-based practice project was to compare mean blood pressures leading up to surgery among patients at Providence Sacred Heart Medical Center (PSHMC).

Methods

- Literature review identified five high quality studies to inform this project.
- Institutional approval sought and obtained. IRB determined exempt research
- Data on adult, elective surgical patients receiving a general anesthetic from 09/30/2017-10/1/2018 were extracted in an anonymized fashion.
- Patients characterized by demographic and clinical characteristics (table 1).
- Mean blood pressures calculated from three settings: ambulatory clinics in the 12 months prior to surgery, the surgical admit unit (SAU) and the "first on table" (FOT) or first pressure recorded in the operating room. Patients without blood pressures recorded in each care setting excluded
- Repeated measures ANOVA and multiple linear regression (α=0.05).



Sacred Heart Medical Center & Children's Hospital

Providence Sacred Heart Medical Center Gonzaga University Nurse Anesthesia Program

School of Nursing & Human Physiology

Fi	nd	lin	gs

Table 1. Demographic and Clinical Characteristics (N=6,081)					
Characteristic	Values	Count	Percent		
Sex					
	Male	2851	47%		
	Female	3230	53%		
ASA Score					
		223	4%		
	II	2491	41%		
	III	2913	48%		
	IV	196	3%		
	Blank	258	4%		
Service Line					
	Orthopedics	1651	27%		
	Neurosurgery	1102	18%		
	General	959	16%		
	Urology	558	9%		
	Vascular	492	6%		
	ENT	382	6%		
	Gynecology	344	6%		
	Pulmonary	192	3%		
	Gastroenterology	144	2%		
	Plastics	110	2%		
	Dental	70	1%		
	Gynecology Oncology	39	1%		
	Ophthalmology	38	1%		
Comorbidities					
	Hypertension	3236	53%		
	Cerebrovascular Disease	1523	25%		
	PVD	1040	17%		
	Diabetes Mellitus	1025	17%		
	Coronary Artery Disease	972	16%		
	COPD	732	12%		
	Chronic Kidney Disease	569	9%		
	Atrial Fibrillation	442	7%		
	Congestive Heart Failure	289	5%		
	Liver Disease	234	4%		
Characteristic		Mean	SD		
Age (years)		60	16		
BMI (kg/m ²)		30	7		

ASA= American Society of Anesthesiologists; ENT= Ear, Nose, Throat; PVD= Peripheral Vascular Disease; COPD= Chronic Obstructive Pulmonary Disease; BMI= Body Mass Index.

Findings (cont.)

Figure 1. Comparison of Mean Blood Pressures by Care Setting



*Calculated by repeated measures ANOVA. † Mean mmHg +/- SD

Table 2. Independent Factors Predicting FOT SBP

Factor	β	SE	95% CI	P-Value
Age	0.27	0.02	0.23-0.31	<0.001
Sex	1.99	0.57	0.87-3.10	<0.001
ASA Score	-0.67	0.54	-1.73-0.39	0.22
BMI	0.06	0.04	-0.02-0.14	0.16
COPD	-0.63	0.91	-2.41-1.15	0.49
Heart Failure	3.41	1.43	-0.61-6.21	0.02
Atrial Fibrillation	3.14	1.13	0.92-5.36	0.01
Diabetes Mellitus	1.05	0.79	-0.51-2.60	0.19
Hypertension	1.92	0.67	0.61-3.23	0.01
CAD	0.01	0.86	-1.68-1.70	0.99
CKD	2.27	1.02	0.27-4.28	0.03
Liver disease	-2.47	1.47	-5.35-0.40	0.09
CVA	0.59	0.66	-0.70-1.89	0.37
PVD	2.17	0.86	0.48-3.85	0.02

 $R^2 = 0.0634$

ASA= American Society of Anesthesiologists; BMI= Body Mass Index; COPD= Chronic Obstructive Pulmonary Disease; CAD= Coronary Artery Disease; CKD= Chronic Kidney Disease; CVA= Cerebrovascular Accident; PVD= Peripheral Vascular Disease

SBP increased significantly from the pre-surgical ambulatory period to the first on table (FOT) blood pressure taken in the operating room. Changes in MAP were statistically, but not clinically, significant. A multivariable regression model revealed age, sex, heart failure, atrial fibrillation, hypertension, chronic kidney disease and peripheral vascular disease to be factors predictive of increasing FOT SBP. Our findings are consistent with current research evidence.

Providence Sacred Heart Medical Center patients may experience over-treatment for hypotension if the first-ontable SBP is utilized as a baseline. This overtreatment may result in increased monetary and physical costs.

Cheung, A. T. (2006). Exploring an optimum intra/postoperative management strategy for acute hypertension in the cardiac surgery patient. Journal of Cardiac Surgery, 21(SUPPL. 1). https://doi.org/10.1111/j.1540-8191.2006.00214.x

Drummond, J. C., Blake, J. L., Patel, P. M., Clopton, P., & Schulteis, G. (2013). An observational study of the influence of "white-coat hypertension" on day-of-surgery blood pressure determinations. Journal of *Neurosurgical Anesthesiology*, *25*(2), 154–161. https://doi.org/10.1097/ANA.0b013e31827a0151

Meng, L., Yu, W., Wang, T., Zhang, L., Heerdt, P. M., & Gelb, A. W. (2018). Blood Pressure Targets in Perioperative Care. Hypertension (Dallas, Tex.: 1979), 72(4), 806–817. https://doi.org/10.1161/HYPERTENSIONAHA.118.11688

Mizuno, J., Kato, S., Sato, T., & Sawamura, S. (2012). Pre-anesthesia systolic blood pressure increases with age regardless of sex. Journal of Anesthesia, 26(4), 496–502. https://doi.org/10.1007/s00540-012-1379-5

Schonberger, R. B., Burg, M. M., Holt, N., Lukens, C. L., Dai, F., & Brandt, C. (2012). The relationship between preoperative and primary care blood pressure among veterans presenting from home for surgery: Is there evidence for anesthesiologist-initiated blood pressure referral? Anesthesia and Analgesia, 114(1), 205–214. https://doi.org/10.1213/ANE.0b013e318239c4c1

Soo, J. C. L., Lacey, S., Kluger, R., & Silbert, B. S. (2011). Defining intra-operative hypotension - a pilot comparison of blood pressure during sleep and general anaesthesia*. *Anaesthesia*, 66(5), 354–360. https://doi.org/10.1111/j.1365-2044.2011.06657.x

Discussion

References

Ard, J. L., & Kendale, S. (2016). Searching for baseline blood pressure: A comparison of blood pressure at three different care points. Journal of Clinical Neuroscience, 34, 59-62. https://doi.org/10.1016/j.jocn.2016.08.001

Bartels, K., Esper, S. A., & Thiele, R. H. (2016). Blood pressure monitoring for the anesthesiologist: A practical review. Anesthesia and Analgesia, 122(6), 1866–1879. https://doi.org/10.1213/ANE.00000000001340